

## AQ 10 WG

### COMPATIBILITY CHART

### FUNGICIDES

Aug. 2013

ACTIVE INGREDIENT	TRADE NAME	COMPATIBILITY
AZOXYSTROBIN	QUADRI	
<i>Bacillus subtilis</i> strain QST713	SERENADE MAX	
<i>Bacillus amyloliquefaciens</i> strain D747	AMYLO-X	
BENALAXYL	various	
BITERTANOLE	PROCLAIM	
BORDEAUX MIXTURE	Copper 20%	
BOSCALID	CANTUS	
BUPIRIMATE	NIMROD	-
CAPTAN	various	
CHLOROTHALONIL	DACONIL	
COPPER HYDROXIDE	KOCIDE/HELIOCUIVRE	
COPPER OXYCHLORIDE	various	
CYAZOFAMID	MILDICUT	
CYFLUFENAMID	TAKUMI	
CYMOXANIL	CURZATE 50%	
CYPROCONAZOLE	ATEMI/CADDY	
CYPRODINIL + FLUDIOXONIL	SWITCH	
DIFENCONAZOLO	SCORE	-
DIMETOMORPH	FORUM 50 WP	
DITHIANON	DELAN	
DODINE	SYLLIT	
FAMOXADONE	EQUATION PRO	
FENAMIDONE	ELICIO	
FENBUCONAZOLE	INDAR	-
FENHEXAMIDE	TELDOR	
FENPYRAZAMINE	PROLECTUS	
FLUAZINAM	OHAYO	
FLUOPYRAM	LUNA PRIVILEGE	
FOLPET	various	
FOSETYL-ALUMINIUM	ALIETTE 80%	
IMAZALIL	DECCOZIL	-
IPRODIONE	ROVRAL	
IPROVALICARB	MELODY COMPACT	
KRESOXIM METHYL	STROBY	
LIME SULPHUR	POLICALCIO	-
MANCOZEB	DITHANE DG	
MANDIPROPAMID	PERGADO SC	
MANEB	various	

MEPANIPYRIM	FRUPICA	
MEPTYLDINOCAP	KARATHANE STAR	
METALAXYL	RIDOMIL	
METRAFENONE	VIVANDO	
METIRAM	POLYRAM DF	
MYCLOBUTANIL	THIOCUR	
PENCONAZOLE	TOPAS	
POTASSIUM BICARBONATE	VITISAN	
PROCHLORAZ	various	-
PROPAMOCARB	PREVICUR	-
PROPICONAZOLE	TILT	
PROPINEB	ANTRACOL	
PROQUINAZID	TALENDO	
PYRIMETHANIL	SCALA	
QUINOXYFEN	ARIUS	
SPIROXAMINE	PROSPER	
SULPHUR	various	
SULPHUR + TERPENIC ALCOHOL	HELIOSOUFRE S	
TEBUCONAZOLE	FOLICUR	
TETRACONAZOLE	DOMARK	-
THIOPHANATE-METHYL	various	
THIABENDAZOLE	various	-
THIRAM	various	
TOLCLOFOS-METHYL	various	-
TRIADIMENOL	BAYFIDAN	
TRIFLOXYSTROBIN	FLINT	
ZIRAM	various	
ZOXAMIDE	ELECTIS R	

### EXPLANATORY NOTES

	<b>COMPATIBLE in tank mixture</b> without any significant reduction in spore germination
	<b>PARTIALLY COMPATIBLE</b> , a certain level of reduction in spore germination could occur; in case of tank mix with AQ 10, apply spray solution within 30 minutes from tank mixing
	<b>NOT COMPATIBLE, do not tank mix with AQ 10.</b> Apply AQ 10 separately and keep a spray interval of at least <b>5 days</b> between the fungicide and AQ 10 treatment
-	<b>NO DATA AVAILABLE</b>

**N.B.** All products were tested at the recommended field rates according to label instructions. Tank mix AQ 10 with fungicides only at recommended label rates. Never mix AQ 10 with a fungicide in its concentrated form. The trade name of the fungicide is provided in order to provide indications on the tested formulation of the active substance. Other formulations may be more or less harmful. In case 2 or more fungicides are tank mixed, consider always the worst case. This edition is for international purposes, some products may not be registered in some countries or on certain crops. No liability will be accepted for any loss or damage as a result of using products listed in this chart.

For any further information please contact:



**BIOGARD – TECHNICAL AREA, Division of CBC (Europe) S.r.l.**

Via Calcinaro 2085/ int.7, 47521 Cesena (FC), Italy

Phone. +39 0547 630336 – Fax +39 0547 632685

e-mail: [tecnicobiogard@cbceurope.it](mailto:tecnicobiogard@cbceurope.it)